

Beers Criteria for Potentially Inappropriate Medication Use in Older Adults Part I: 2002 Criteria Independent of Diagnoses or Conditions

By: Sheila Molony, PhD, RN, BC, GNP, Yale University School of Nursing

WHY: Recently published studies confirm that inappropriate medication use remains a serious problem for the elderly (Bonk, et al, 2006; Lau, et al, 2005). Nursing knowledge of potentially inappropriate medications will enable attentive monitoring for adverse effects, and increase awareness of disease and condition-specific medication concerns in older adults.

BEST TOOL: The 2002 *Criteria for Potentially Inappropriate Medication Use in Older Adults* (Fick, et al, 2003) update the 1997 Beers Criteria, and identify medications noted by an expert consensus panel to have potential risks that outweigh potential benefits. The criteria provide an outcome severity rating (high vs. low) and a brief summary of the prescribing concerns. A total of 15 medications/classes were dropped or modified from the 1997 list, and 44 new medications were added.

Try This: Part I focuses on Table 1: 2002 Criteria Independent of Diagnoses and Conditions. It contains 48 individual medications or classes of medications to avoid in older adults and their potential concerns. Try This: Part II presents the criteria considering diagnosis or medical condition.

TARGET POPULATION: The criteria apply to the general population of adults older than 65 years of age. There may be additional medications that are inappropriate for a significantly older or frailer population.

VALIDITY AND RELIABILITY: The criteria were developed using a modified Delphi method to achieve consensus among 12 experts in geriatrics and/or pharmacology. The criteria have been used to screen populations for possible medication-related problems. Use of inappropriate medications has been associated with negative outcomes (Fu, et al, 2004; Lau, et al, 2005; Perri, et al, 2005). Additional studies are needed to support predictive validity and address potential confounding variables such as severity of underlying illness (Lau, et al, 2005; Zuckerman, et al, 2006).

STRENGTHS AND LIMITATIONS: The criteria will assist nurses to identify patients who may benefit from monitoring or medication review. The criteria do not identify all cases of potentially inappropriate prescribing or medication-associated adverse events, and do not address polypharmacy or underuse of helpful medications. The criteria are designed for population-based screening and are not intended to substitute for professional judgment regarding the individualized needs of particular older adults.

FOLLOW-UP: Nurses may use the criteria to increase awareness of medications that may increase risk for adverse drug reactions. Nurses, primary care providers and pharmacists may collaborate to optimize individualized medication regimes and provide appropriate clinical monitoring and education. The suggested references provide further information on medication risk and older adults.

MORE ON THE TOPIC:

Best practice information on care of older adults: www.ConsultGerIRN.org.

Beers, M.H. (1997). Explicit criteria for determining potentially inappropriate medication use by the elderly.

Archives of Internal Medicine, 157, 1531-1536.

Bonk, M.E., Krown, H., Matuszewski, K., & Oinonen, M. (2006). Potentially inappropriate medications in hospitalized senior patients. *American Journal of Health-System Pharmacy*, 63(12), 1161-1165.

Fick, D.M., Cooper, J.W., Wade, W.E., Waller, J.L., Maclean, J.R., & Beers, M.H. (2003). Updating the Beers Criteria for potentially inappropriate medication use in older adults: Results of a US consensus panel of experts. *Archives of Internal Medicine*, 163(22), 2716-2724.

Fu, A.Z., Liu, G.G., & Christensen, D.B. (2004). Inappropriate medication use and health outcomes in the elderly. *JAGS*, 52(11), 1934-1939.

Lau, D.T., Kasper, J.D., Potter, D.E., Lyles, A., & Bennett, R.G. (2005). Hospitalization and death associated with potentially inappropriate medication prescriptions among elderly nursing home residents. *Archives of Internal Medicine*, 165(1), 68-74.

Perri, M. III, Menon, A.M., Deshpande, A.D., Shinde, S.B., Jiang, R., & Cooper, J.W., Cook, C.L., Griffin, S.C., & Lorys, R.A. (2005). Adverse outcomes associated with inappropriate drug use in nursing homes. *Annals of Pharmacotherapy*, 39(3), 405-411.

Zuckerman, I.H., Langenberg, P., Baumgarten, M., Orwig, D., Byrns, P.J., & Simoni-Wastila, L., & Magaziner, J. (2006).

Inappropriate drug use and risk of transition to nursing homes among community-dwelling older adults. *Medical care*, 44(8), 722-730.

Table 1: 2002 Criteria for Potentially Inappropriate Medication Use in Older Adults: Independent of Diagnoses or Conditions

Drug	Concern	Severity Rating (High or Low)
Propoxyphene (Darvon) and combination products (Darvon with ASA, Darvon-N, and Darvocet-N)	Offers few analgesic advantages over acetaminophen, yet has the adverse effects of other narcotic drugs.	Low
Indomethacin (Indocin and Indocin SR)	Of all available nonsteroidal anti-inflammatory drugs, this drug produces the most CNS adverse effects.	High
Pentazocine (Talwin)	Narcotic analgesic that causes more CNS adverse effects, including confusion and hallucinations, more commonly than other narcotic drugs. Additionally, it is a mixed agonist and antagonist.	High
Trimethobenzamide (Tigan)	One of the least effective antiemetic drugs, yet it can cause extrapyramidal adverse effects.	High
Muscle relaxants and antispasmodics: methocarbamol (Robaxin), carisoprodol (Soma), chlorzoxazone (Paraflex), metaxalone (Skelaxin), cyclobenzaprine (Flexeril), and oxybutynin (Ditropan). Do not consider the extended-release Ditropan XL.	Most muscle relaxants and antispasmodic drugs are poorly tolerated by elderly patients, since these cause anticholinergic adverse effects, sedation, and weakness. Additionally, their effectiveness at doses tolerated by elderly patients is questionable.	High
Flurazepam (Dalmane)	This benzodiazepine hypnotic has an extremely long half-life in elderly patients (often days), producing prolonged sedation and increasing the incidence of falls and fracture. Medium- or short-acting benzodiazepines are preferable.	High
Amitriptyline (Elavil), chlordiazepoxide-amitriptyline (Limbital), and perphenazine-amitriptyline (Triavil)	Because of its strong anticholinergic and sedation properties, amitriptyline is rarely the antidepressant of choice for elderly patients.	High
Doxepin (Sinequan)	Because of its strong anticholinergic and sedating properties, doxepin is rarely the antidepressant of choice for elderly patients.	High
Meprobamate (Miltown and Equanil)	This is a highly addictive and sedating anxiolytic. Those using meprobamate for prolonged periods may become addicted and may need to be withdrawn slowly.	High
Doses of short-acting benzodiazepines: doses greater than lorazepam (Ativan), 3 mg; oxazepam (Serax), 60 mg; alprazolam (Xanax), 2 mg; temazepam (Restoril), 15 mg; and triazolam (Halcion), 0.25 mg	Because of increased sensitivity to benzodiazepines in elderly patients, smaller doses may be effective as well as safer. Total daily doses should rarely exceed the suggested maximums.	High
Long-acting benzodiazepines: chlordiazepoxide (Librium), chlordiazepoxide-amitriptyline (Limbital) clidinium-chlordiazepoxide (Librax), diazepam (Valium), quazepam (Doral), halazepam (Paxipam), and chlorazepate (Tranxene)	These drugs have a long half-life in elderly patients (often several days), producing prolonged sedation and increasing the risk of falls and fractures. Short- and intermediate-acting benzodiazepines are preferred if a benzodiazepine is required.	High
Disopyramide (Norpace and Norpace CR)	Of all antiarrhythmic drugs, this is the most potent negative inotrope and therefore may induce heart failure in elderly patients. It is also strongly anticholinergic. Other antiarrhythmic drugs should be used.	High
Digoxin (Lanoxin) (should not exceed >0.125 mg/d except when treating atrial arrhythmias)	Decreased renal clearance may lead to increased risk of toxic effects.	Low
Short-acting dipyrindamole (Persantine). Do not consider the long-acting dipyrindamole (which has better properties than the short-acting in older adults) except with patients with artificial heart valves	May cause orthostatic hypotension.	Low
Methyldopa (Aldomet) and methyldopa-hydrochlorothiazide (Aldoril)	May cause bradycardia and exacerbate depression in elderly patients.	High
Reserpine at doses >0.25 mg	May induce depression, impotence, sedation, and orthostatic hypotension.	Low
Chlorpropamide (Diabinese)	It has a prolonged half-life in elderly patients and could cause prolonged hypoglycemia. Additionally, it is the only oral hypoglycemic agent that causes SIADH.	High
Gastrointestinal antispasmodic drugs: dicyclomine (Bentyl), hyoscyamine (Levsin and Levsinex), propantheline (Pro-Banthine), belladonna alkaloids (Donnatal and others), and clidinium-chlordiazepoxide (Librax)	GI antispasmodic drugs are highly anticholinergic and have uncertain effectiveness. These drugs should be avoided (especially for long-term use).	High
Anticholinergics and antihistamines: chlorpheniramine (Chlor-Trimeton), diphenhydramine (Benadryl), hydroxyzine (Vistaril and Atarax), cyproheptadine (Pericatin), promethazine (Phenergan), tripeleminamine, dexchlorpheniramine (Polaramine)	All nonprescription and many prescription antihistamines may have potent anticholinergic properties. Nonanticholinergic antihistamines are preferred in elderly patients when treating allergic reactions.	High
Diphenhydramine (Benadryl)	May cause confusion and sedation. Should not be used as a hypnotic, and when used to treat emergency allergic reactions, it should be used in the smallest possible dose.	High
Ergot mesyloids (Hydergine) and cyandelate (Cyclospasmol)	Have not been shown to be effective in the doses studied.	Low
Ferrous sulfate >325 mg/d	Doses >325 mg/d do not dramatically increase the amount absorbed but greatly increase the incidence of constipation.	Low
All barbiturates (except phenobarbital) except when used to control seizures	Are highly addictive and cause more adverse effects than most sedative or hypnotic drugs in elderly patients.	High
Meperidine (Demerol)	Not an effective oral analgesic in doses commonly used. May cause confusion and has many disadvantages to other narcotic drugs.	High
Ticlopidine (Ticlid)	Has been shown to be no better than aspirin in preventing clotting and may be considerably more toxic. Safer, more effective alternatives exist.	High
Ketorolac (Toradol)	Immediate and long-term use should be avoided in older persons, since a significant number have asymptomatic GI pathologic conditions.	High
Amphetamines and anorexic agents	These drugs have potential for causing dependence, hypertension, angina, and myocardial infarction.	High
Long-term use of full-dosage, longer half-life, non-COX-selective NSAIDs: naproxen (Naprosyn, Avapro, and Aleve), oxaprozin (Daypro), and piroxicam (Feldene)	Have the potential to produce GI bleeding, renal failure, high blood pressure, and heart failure.	High
Daily fluoxetine (Prozac)	Long half-life of drug and risk of producing excessive CNS stimulation, sleep disturbances, and increasing agitation. Safer alternatives exist.	High
Long-term use of stimulant laxatives: bisacodyl (Dulcolax), cascara sagrada, and Neoloid except in the presence of opiate analgesic use	May exacerbate bowel dysfunction.	High
Amiodarone (Cordarone)	Associated with QT interval problems and risk of provoking torsades de pointes. Lack of efficacy in older adults.	High
Orphenadrine (Norflex)	Causes more sedation and anticholinergic adverse effects than safer alternatives.	High
Guanethidine (Ismelin)	May cause orthostatic hypotension. Safer alternatives exist.	High
Guanadrel (Hylorel)	May cause orthostatic hypotension.	High
Cyclandelate (Cyclospasmol)	Lack of efficacy.	Low
Isoxsuprine (Vasodilan)	Lack of efficacy.	Low
Nitrofurantoin (Macrochantin)	Potential for renal impairment. Safer alternatives available.	High
Doxazosin (Cardura)	Potential for hypotension, dry mouth, and urinary problems.	Low
Methyltestosterone (Android, Virilon, and Testrad)	Potential for prostatic hypertrophy and cardiac problems.	High
Thioridazine (Mellaril)	Greater potential for CNS and extrapyramidal adverse effects.	High
Mesoridazine (Sereniti)	CNS and extrapyramidal adverse effects.	High
Short acting nifedipine (Procardia and Adalat)	Potential for hypotension and constipation.	High
Clonidine (Catapres)	Potential for orthostatic hypotension and CNS adverse effects.	Lo
Mineral oil	Potential for aspiration and adverse effects. Safer alternatives available.	High
Cimetidine (Tagamet)	CNS adverse effects including confusion.	Low
Ethacrynic acid (Edecrin)	Potential for hypertension and fluid imbalances. Safer alternatives available.	Low
Desiccated thyroid	Concerns about cardiac effects. Safer alternatives available.	High
Amphetamines (excluding methylphenidate hydrochloride and anorexics)	CNS stimulant adverse effects.	High
Estrogens only (oral)	Evidence of the carcinogenic (breast and endometrial cancer) potential of these agents and lack of cardioprotective effect in older women.	Low

Abbreviations: CNS, central nervous system; COX, cyclooxygenase; GI, gastrointestinal; NSAIDs, nonsteroidal anti-inflammatory drugs; SIADH, syndrome of inappropriate antidiuretic hormone secretion. Reprinted with permission: Fick, D.M., Cooper, J.W., Wade, W.E., Waller, J.L., Maclean, J.R., & Beers, M.H. (2003). Updating the Beers Criteria for potentially inappropriate medication use in older adults: Results of a US consensus panel of experts. Archives of Internal Medicine, 163(22), 2716-2724. Table 1, p. 2720. Evidence Level VI: Expert Opinion. Copyright © 2003, American Medical Association. All Rights reserved.



A SERIES PROVIDED BY
The Hartford Institute for Geriatric Nursing
 EMAIL: hartford.ign@nyu.edu
 HARTFORD INSTITUTE WEBSITE: www.hartfordign.org
 CONSULTGERIERN WEBSITE: www.ConsultGerIERN.org